arXiv:1304.7203v1 [math-ph] 26 Apr 2013

A method for computing the generating function of the characters of simple Lie algebras

J. Fernández Núñez[†], W. García Fuertes[†], A.M. Perelomov[‡]

[†] *Dep. de Física, Facultad de Ciencias, Universidad de Oviedo, Spain*

[‡] *Institute for Theoretical and Experimental Physics, Moscow, Russia.*

Abstract

We describe a general method, based on elementary considerations, to obtain the generating functions of the characters of simple Lie algebras. We show how the method works by means of a few examples involving some low rank classical algebras. As an application, we compute several generating functions for the weight multiplicities of the irreducible representations of these algebras.

PACS: 02.20.Qs, 02.30.Ik, 03.65.Fd
MCS: 17B10, 17B80, 81R12

Key words: Lie algebras, representation theory, quantum integrable systems

1 Description of the method

The characters of the irreducible representations of the simple Lie algebras are systems of orthogonal polynomials which enjoy many interesting properties and have a distinguised role in pure mathematics and mathematical physics [1]. Many features of orthogonal systems of this kind can be studied by means of their generating functions, defined as formal power series in some auxiliary variables whose coefficients give the polynomials entering in the system. In the case of the characters of a simple Lie algebra, the most usual approach for the computation of the generating functions is based on the Weyl character formula [2, 3] and requires, in consequence, to perform a sum over the elements of the Weyl group which is, in general, a rather cumbersome combinatorial task. While this strategy is powerful and can be used to establish formulas of great generality, see for instance [4, 5], we feel that it would be also desirable to put forward an alternative approach which treats the problem in a more elementary fashion, being at the same time able to yield the generating functions of the characters of a particular algebra though a few simple steps.

Thus, in this paper, we take this point of view and put the focus not on general abstract formulas but, instead, in a concrete and quite versatile algorithm which is suitable to be applied separately to each particular algebra. The possibility of such an algorithm comes from two facts. First, in the last few years it has been shown explicitly how the quantum theory of integrable systems, in particular that of the Calogero-Sutherland model [6, 7, 8, 9, 10], can be used to compute the characters of the simple Lie algebras by identifying them with the eigenfunctions of the Hamiltonian for some particular values of the coupling constants, and several lists of characters obtained from this method are now available [11, 12, 13]. And, second, although the procedure that we are going to develop involves some long calculations, this is not a serious drawback, because they are quite straightforward and can be performed rather quickly by means of symbolic calculus languages like Mathematica or Maple, nowadays of common use. In this respect, we remark that, although we shall illustrate here the approach by applying it only to some low-rank algebras, the use of these programs turns the method useful also for higher rank ones.

Let $\mathcal{A}$ be a simple Lie algebra of rank r with fundamental weights $\lambda_1, \lambda_2, \ldots, \lambda_r$ and let us denote R_λ the irreducible representation of $\mathcal{A}$ with highest weight $\lambda = m_1\lambda_1 + m_2\lambda_2 + \cdots + m_r\lambda_r$. The character of this representation is defined as

$$\boldsymbol{\chi}_{m_1,m_2,\ldots,m_r} = \sum_w n_w e(w) \tag{1}$$

where the sum extends to all weights w entering in the representation, n_w is the mutiplicity of the weight w and, if $w = n_1\lambda_1 + n_2\lambda_2 + \cdots + n_r\lambda_r$, then $e(w)$ is

$$e(w) = \exp\left(i\sum_{l=1}^{r} n_l\varphi_l\right) = x_1^{n_1}x_2^{n_2}\cdots x_r^{n_r}, \tag{2}$$

where $\varphi_1, \varphi_2, \ldots, \varphi_r$ are angular coordinates on the maximal torus and x_l are complex phases, $x_l = e^{i\varphi_l}$. In particular, we will denote the character of the representations R_{λ_k} for $k = 1, 2, \ldots, r$ as z_k and we will write all the characters as polynomials in the r variables z_k.

The generating function for the characters of $\mathcal{A}$ is the formal series

$$G(t_j; z_k) = \sum_{m_1=0}^{\infty}\sum_{m_2=0}^{\infty}\cdots\sum_{m_r=0}^{\infty} t_1^{m_1}t_2^{m_2}\cdots t_r^{m_r}\boldsymbol{\chi}_{m_1,m_2,\ldots,m_r}(z_1, z_2, \ldots, z_r) \tag{3}$$

and we expect it to be a rational function [2, 3], namely

$$G(t_j; z_k) = \frac{N(t_j; z_k)}{D(t_j; z_k)}$$

where both the numerator $N(t_j; z_k)$ and the denominator $D(t_j; z_k)$ are polynomials in the t-variables with coefficients depending on the z-variables. We now describe how to compute $D(t_j; z_k)$ and $N(t_j; z_k)$ along four sucessive steps:

1. The term $t_1^{m_1}t_2^{m_2}\cdots t_r^{m_r}\boldsymbol{\chi}_{m_1,m_2,\ldots,m_r}$ of the generating function comes from the representation $R_{m_1\lambda_1+m_2\lambda_2+\cdots+m_r\lambda_r}$, which is part of the tensor product $R_{\lambda_1}^{m_1} \otimes R_{\lambda_2}^{m_2} \otimes \cdots \otimes R_{\lambda_r}^{m_r}$. This implies that the character of that representation includes a term of the form $\prod_{j=1}^{r} e(w_j)^{m_j}$ for each possible combination formed by picking up a weight w_j of each R_{λ_j}. We can obtain all these products multiplied by $t_1^{m_1}t_2^{m_2}\cdots t_r^{m_r}$ if we multiply the geometric series $\sum_{p=0}^{\infty}(t_j)^p e(w)^p$ for all

nonzero weights w entering in the representations of each R_{λ_j}. Therefore, the denominator of the generating function has the general form

$$D = D_1 \times D_2 \times \cdots \times D_r\,, \tag{4}$$

with the factor D_j related to the fundamental representation R_{λ_j} through the formula

$$D_j = \prod_w (1 - t_j^n e(w)) \tag{5}$$

where n is the multiplicity of w in R_{λ_j}, and the product is extended to all non-zero weights of R_{λ_j}. Expanding this product, one can write each D_j as a polynomial in t_j with coefficients depending on the phases x_l but, as the characters are invariant under Weyl reflections, so are the coefficients of these polynomials. As a consequence, they can be rewritten in terms of the Weyl invariant variables z_k. So, in this first step we compute each D_j by means of (5) and use (1) and (2) to put it as a function of the z-variables. Finally, (4) yields $D(t_j; z_k)$.

2. In the second step we compute the generating function

$$F(t_1, t_2, \ldots, t_r) = \sum_{m_1=0}^{\infty} \sum_{m_2=0}^{\infty} \cdots \sum_{m_r=0}^{\infty} t_1^{m_1} t_2^{m_2} \cdots t_r^{m_r} \dim R_{m_1\lambda_1 + m_2\lambda_2 + \cdots + m_r\lambda_r} \tag{6}$$

for the dimensions of the ireducible representations of the algebra $\mathcal{A}$. This generating function is related to $G(t_j; z_k)$ through the replacement $z_k \to \dim R_{\lambda_k}$, i.e. $F(t_j) = G(t_j; \dim R_{\lambda_k})$, where we can compute the dimension of any desired representation by means of the Weyl formula. Let us write

$$F(t_j) = \frac{P(t_j)}{Q(t_j)}. \tag{7}$$

The denominator is known, $Q(t_j) = D(t_j; \dim R_{\lambda_k})$, and for the numerator we will try an ansatz of the form

$$P(t_1, t_2, \ldots, t_r) = \sum_{l_1=0}^{d_1} \sum_{l_2=0}^{d_2} \cdots \sum_{l_r=0}^{d_r} t_1^{l_1} t_2^{l_2} \cdots t_r^{l_r} p_{l_1, l_2, \ldots, l_r}\,, \tag{8}$$

where d_j is the number of non-zero weights in R_{λ_j}, $p_{0,0,\ldots,0} = 1$ and the remaining mumerical coefficientes $p_{l_1, l_2, \ldots, l_r}$ are to be determined by comparing the Taylor expansion of (7) with the right member of (6). It will turn out that most of these coeffients are zero and the final form of $F(t_j)$ can be fixed after a few steps involving some very simple equations.

3. In the third step we come back to the numerator $N(t_j; z_k)$ through the ansatz

$$N(t_j; z_k) = \sum_{l_1, l_2, \ldots, l_r} C_{l_1, l_2, \ldots, l_r}(z_k) t_1^{l_1} t_2^{l_2} \ldots t_r^{l_r}\,,$$

where we include only the terms corresponding to the non-vanishing $p_{l_1, l_2, \ldots, l_r}$ coefficients. Often, we can guess some of the $C_{l_1, l_2, \ldots, l_r}(z_k)$ directly from the numerical value of the coefficient $p_{l_1, l_2, \ldots, l_r}$. For the others, we will have to write explicitly some low order terms of the right member of (3) and compare them with the Taylor expansion of $G(t_j; z_k)$. Once this is done, we have a tentative expression for $G(t_j; z_k)$.

4. The remaining step is to check that our tentative generating function is indeed correct. To this end, we recall that the Schrödinger equation for the Calogero- erland model related to the Lie algebra $\mathcal{A}$ leads, for some specific values of the coupling constants, to an eigenvalue problem of the form
$$\Delta_z \Phi_{m_1,m_2,\ldots,m_r}(z_k) = \varepsilon(m_1, m_1, \ldots, m_r)\Phi_{m_1,m_2,\ldots,m_r}(z_k)$$
where Δ_z is a second order differential operator in the variables z_k, the eigenvalues are given by a known expression involving the inner product in the weight-space of the algebra and the egenfunctions are precisely the characters of $\mathcal{A}$,
$$\Phi_{m_1,m_2,\ldots,m_r}(z_k) = \boldsymbol{\chi}_{m_1,m_2,\ldots,m_r}(z_k),$$
see the references [10, 11, 12, 13] for the details. So, due to the definition (3), the generating function for characters is a solution of the differential equation
$$(\Delta_t - \Delta_z)G(t_j; z_k) = 0\,, \tag{9}$$
with Δ_t being the differential operator
$$\Delta_t = \varepsilon(t_1\partial_{t_1}, t_2\partial_{t_2}, \ldots, t_r\partial_{t_r}).$$
By verifying that our tentative expression satisfies (9), we can finally prove that it gives correctly the generating function we are looking for.

2 Some examples

Let us now apply the method described in the previous section to some Lie algebras in order to appreciate how it works. We will work out the cases of the simple classical Lie algebras up to rank two, i.e. $A_1 = B_1 = C_1 = D_1$, A_2 and $C_2 \simeq B_2$, since $D_2 \simeq A_1 \oplus A_1$ is semisimple.

2.1 The generating function for the algebra A_1

This is the simplest case. The algebra has only a fundamental weight λ_1 and the fundamental representation R_{λ_1} has weights $\{\lambda_1, -\lambda_1\}$. Thus, the character z_1 is
$$z_1 = e^{i\varphi_1} + e^{-i\varphi_1} = x_1 + \frac{1}{x_1} \tag{10}$$
The generating functions for characters and dimensions are
$$G(t_1; z_1) = \sum_{m_1=0}^{\infty} t_1^{m_1}\boldsymbol{\chi}_{m_1}(z_1)\,, \qquad F(t_1) = \sum_{m_1=0}^{\infty} t_1^{m_1} \dim R_{m_1}\,.$$
Let us now apply our four-step method. As there only a weight, the denominator $D(t_1; z_1)$ is simply $D(t_1; z_1) = D_1$ with
$$D_1 = (1 - t_1x_1)(1 - t_1\frac{1}{x_1}) = 1 - t_1z_1 + t_1^2\,.$$

Now, as the dimension of R_1 is two, the denominator of the generating function for dimensions is $Q(t_1) = D(t_1; 2) = (1 - t_1)^2$. For the numerator, we try the ansatz $P(t_1) = 1 + p_1 t_1 + p_2 t_1^2$. The Weyl formula for dimensions gives $\dim R_{m_1} = m_1 + 1$ and comparing this ansatz with the first few terms of the series $F(t_1)$, one readily finds that $p_1 = p_2 = 0$. Therefore

$$F(t_1) = \frac{1}{(1-t)^2}.$$

Given the form of $F(t_1)$, our ansatz for the numerator for generating function for characters is

$$N(t_1; z_1) = C_0(z_1)$$

but, given that $d_0 = 1$, the only sensible guess is $C_0(z_1) = 1$ and the tentative expression for the generating function is

$$G(t_1; z_1) = \frac{1}{1 - t_1 z_1 + t_1^2}. \tag{11}$$

It remains to confirm that this is correct. For that, we resort to the Calogero-Sutherland theory related to the algebra A_1 (see for example[14]), which leads to the differential operator

$$\Delta_z = (z_1^2 - 4)\partial_{z_1}^2 + 3z_1 \partial_{z_1}\,,$$

with eigenvalues given by

$$\varepsilon(m_1) = m_1^2 + 2m_1.$$

Thus, the operator Δ_t is $\Delta_t = t_1^2 \partial_{t_1} + 3t_1 \partial_{t_1}$ and one can check that

$$(\Delta_t - \Delta_z) \frac{1}{1 - t_1 z_1 + t_1^2} = 0\,,$$

as it should be. Therefore (11) gives correctly the generating function for the characters of A_1.

2.2 The generating function for the algebra A_2

In this case there are two fundamental weights λ_1, λ_2, and the fundamental representations R_{λ_1} and R_{λ_2} have, respectively, weights $\{\lambda_1, \lambda_2 - \lambda_1, -\lambda_2\}$ and $\{\lambda_2, \lambda_1 - \lambda_2, -\lambda_1\}$. Therefore, the characters z_1 and z_2 are

$$z_1 = e^{i\varphi_1} + e^{-i\varphi_2} + e^{i(\varphi_2 - \varphi_1)} = x_1 + \frac{1}{x_2} + \frac{x_2}{x_1} \tag{12}$$

$$z_2 = e^{i\varphi_2} + e^{-i\varphi_1} + e^{i(\varphi_1 - \varphi_2)} = x_2 + \frac{1}{x_1} + \frac{x_1}{x_2}. \tag{13}$$

The generating functions to be computed are

$$G(t_1, t_2; z_1, z_2) = \sum_{m_1=0}^{\infty} \sum_{m_2=0}^{\infty} t_1^{m_1} t_2^{m_2} \boldsymbol{\chi}_{m_1,m_2}(z_1, z_2) \tag{14}$$

$$F(t_1, t_2) = \sum_{m_1=0}^{\infty} \sum_{m_2=0}^{\infty} t_1^{m_1} t_2^{m_2} \dim R_{m_1,m_2} \tag{15}$$

and we follow the same steps than before. The denominator of the generating function for characters is $D(t_1,t_2;z_1,z_2) = D_1 \times D_2$ where

$$\begin{aligned} D_1 &= (1-t_1x_1)(1-t_1\frac{1}{x_2})(1-t_1\frac{x_2}{x_1}) = 1-t_1z_1+t_1^2z_2-t_1^3 \\ D_2 &= (1-t_2x_2)(1-t_2\frac{1}{x_1})(1-t_2\frac{x_1}{x_2}) = 1-t_2z_2+t_2^2z_1-t_2^3. \end{aligned}$$

Since both R_{λ_1} and R_{λ_2} have dimension three, the denominator of the generating function for dimensions is $Q(t_1,t_2) = D(t_1,t_2;3,3) = (1-t_1)^3(1-t_2)^3$. For the numerator, we try the ansatz

$$P(t_1,t_2) = \sum_{l_1=0}^{3}\sum_{l_2=0}^{3} p_{l_1,l_2} t_1^{l_1} t_2^{l_2}$$

and, using that $\dim R_{m_1,m_2} = \frac{1}{2}(m_1+1)(m_2+1)(m_1+m_2+2)$ to compare with the right member of (15), we find that the only nonvanishing coefficients are $p_{0,0} = -p_{1,1} = 1$. Thus, the generating function for dimensions is

$$F(t_1,t_2) = \frac{1-t_1t_2}{(1-t_1)^3(1-t_2)^3}.$$

In view of that, the ansatz for the numerator for the generating function for characters is

$$N(t_1,t_2;z_1,z_2) = C_{0,0}(z_1,z_2) + C_{1,1}(z_1,z_2)t_1t_2$$

and the reasonable guess is $C_{0,0}(z_1,z_2) = -C_{1,1}(z_1,z_2) = 1$. One can confirm that this guess is correct by comparing the ansatz with the right member of (14): to do so, the only characters needed

$$\boldsymbol{\chi}_{1,0}(z_1,z_2) = z_1, \quad \boldsymbol{\chi}_{0,1}(z_1,z_2) = z_2, \quad \boldsymbol{\chi}_{1,1}(z_1,z_2) = z_1z_2 - 1$$

. Therefore, the tentative form of the generating function is

$$G(t_1,t_2;z_1,z_2) = \frac{1-t_1t_2}{(1-t_1z_1+t_1^2z_2-t_1^3)(1-t_2z_2+t_2^2z_1-t_2^3)}. \tag{16}$$

Now, the second order differential operator associated to the Calogero-Sutherland model for the algebra A_2 whose eigenfunctions are the characters is [11, 12]

$$\Delta_z = (z_1^2-3z_2)\partial_{z_1}^2 + (z_2^2-3z_1)\partial_{z_2}^2 + (z_1z_2-9)\partial_{z_1}\partial_{z_2} + 4z_1\partial_{z_1} + 4z_2\partial_{z_2}$$

and their eigenvalues are

$$\varepsilon(m_1,m_2) = m_1^2 + m_2^2 + m_1m_2 + 3m_1 + 3m_2.$$

Thus, the operator Δ_t is $\Delta_t = t_1^2\partial_{t_1}^2 + t_2^2\partial_{t_2}^2 + t_1t_2\partial_{t_1}\partial_{t_2} + 4t_1\partial_{t_1} + 4t_2\partial_{t_2}$. One can check that, in fact,

$$(\Delta_t - \Delta_z)\frac{1-t_1t_2}{(1-t_1z_1+t_1^2z_2-t_1^3)(1-t_2z_2+t_2^2z_1-t_2^3)} = 0$$

so that (16) is the correct generating function for the characters of A_2.

2.3 The generating function for the algebra C_2

Again, there are two weights, but now the weights in fundamental representations R_{λ_1} and R_{λ_2} are, respectively, $\{\lambda_1, -\lambda_1, \lambda_1-\lambda_2, \lambda_2-\lambda_1\}$ and $\{0, \lambda_2, -\lambda_2, 2\lambda_1-\lambda_2, -2\lambda_1+\lambda_2\}$. Thus, the characters z_1 and z_2 are

$$\begin{aligned} z_1 &= e^{i\varphi_1}+e^{-i\varphi_1}+e^{i(\varphi_1-\varphi_2)}+e^{i(\varphi_2-\varphi_1)} = x_1+\frac{1}{x_1}+\frac{x_1}{x_2}+\frac{x_2}{x_1} && (17)\\ z_2 &= 1+e^{i\varphi_2}+e^{-i\varphi_2}+e^{i(2\varphi_1-\varphi_2)}+e^{i(\varphi_2-2\varphi_1)} = 1+x_2+\frac{1}{x_2}+\frac{x_1^2}{x_2}+\frac{x_2}{x_1^2}. && (18)\end{aligned}$$

The denominator of the generating function for characters is $D(t_1,t_2;z_1,z_2) = D_1\times D_2$ with

$$\begin{aligned} D_1 &= (1-t_1x_1)(1-t_1\frac{1}{x_1})(1-t_1\frac{x_1}{x_2})(1-t_1\frac{x_2}{x_1}) = 1-t_1z_1+t_1^2(z_2+1)-t_1^3z_1+t_1^4\\ D_2 &= (1-t_2x_2)(1-t_2\frac{1}{x_2})(1-t_2\frac{x_1^2}{x_2})(1-t_2\frac{x_2}{x_1^2}) = 1-t_2(z_2-1)+t_2^2(z_1^2-2z_2)-t_2^3(z_2-1)+t_2^4,\end{aligned}$$

while the denominator of the generating function for dimensions is

$$Q(t_1,t_2) = D(t_1,t_2;4,5) = (1-t_1)^4(1-t_2)^4.$$

For the numerator, we try the ansatz

$$P(t_1,t_2) = \sum_{l_1=0}^{4}\sum_{l_2=0}^{4} p_{l_1,l_2}t_1^{l_1}t_2^{l_2}$$

and, by means of $\dim R_{m_1,m_2} = \frac{1}{6}(m_1+1)(m_2+1)(m_1+m_2+2)(m_1+2m_2+3)$, we find that the only nonvanishing coefficients are $p_{0,0}=p_{0,1}=p_{2,1}=p_{2,2}=1$ and $p_{1,1}=-4$. Thus

$$F(t_1,t_2) = \frac{(1+t_2)(1+t_1^2t_2)-4t_1t_2}{(1-t_1)^4(1-t_2)^4}.$$

Given this result, the ansatz for the numerator of the generating function for characters is

$$N(t_1,t_2;z_1,z_2) = C_{0,0}(z_k)+C_{0,1}(z_k)t_2+C_{1,1}(z_k)t_1t_2+C_{1,2}(z_k)t_1t_2^2+C_{2,2}(z_k)t_1^2t_2^2$$

and the reasonable guess is $C_{0,0}=C_{0,1}=C_{2,1}=C_{2,2}=1$ and $C_{1,1}=-z_1$. Again, one can check that this guess is accurate by comparing with the explicit series of characters: the only characters needed are $\boldsymbol{\chi}_{1,0}, \boldsymbol{\chi}_{0,1}, \boldsymbol{\chi}_{1,1}, \boldsymbol{\chi}_{2,1}$ and $\boldsymbol{\chi}_{2,2}$, which are given in the Appendix. Therefore the tentative form of the generating function is

$$G(t_1,t_2;z_1,z_2) = \frac{1+t_2-z_1t_1t_2+t_1^2t_2+t_1^2t_2^2}{(1-(t_1+t_1^3)z_1+t_1^2(z_2+1)+t_1^4)(1-(t_2+t_2^3)(z_2-1)+t_2^2(z_1^2-2z_2)+t_2^4)}. \tag{19}$$

Now we need the Calogero-Sutherland differential operator for C_2, which is shown in the Appendix to be

$$\Delta_z = (z_1^2-2z_2-6)\partial_{z_1}^2+(2z_2^2-4z_1^2+4z_2-6)\partial_{z_2}^2+(2z_1z_2-10z_1)\partial_{z_1}\partial_{z_2}+5z_1\partial_{z_1}+4z_2\partial_{z_2}.$$

Since their eigenvalues are

$$\varepsilon(m_1, m_2) = m_1^2 + 2m_2^2 + 2m_1m_2 + 4m_1 + 6m_2,$$

we see that the operator Δ_t is $\Delta_t = t_1^2\partial_{t_1}^2 + 2t_2^2\partial_{t_2}^2 + 2t_1t_2\partial_{t_1}\partial_{t_2} + 5t_1\partial_{t_1} + 8t_2\partial_{t_2}$. We now can check that

$$(\Delta_t - \Delta_z)\frac{1 + t_2 - z_1t_1t_2 + t_1^2t_2 + t_1^2t_2^2}{(1 - (t_1 + t_1^3)z_1 + t_1^2(z_2 + 1) + t_1^4)(1 - (t_2 + t_2^3)(z_2 - 1) + t_2^2(z_1^2 - 2z_2) + t_2^4)} = 0$$

and this proves that (19) is the correct generating function for the characters of C_2.

2.4 Generating functions for some subsets of characters

If in the previous examples for A_2 and C_2 we take $t_1 = 0$ or $t_2 = 0$, we obtain the generating function for the characters of the form $\boldsymbol{\chi}_{m_1,0}(z_1, z_2)$ or $\boldsymbol{\chi}_{0,m_2}(z_1, z_2)$ of these algebras. It is also possible to use the method that we have been describing to compute the generating function of some other subsets of characters. Let us, for instance, consider the case of the diagonal characters of A_2 and seek for their generating function

$$G_{diag}(t; z_1, z_2) = \sum_{m=0}^{\infty} t^m \boldsymbol{\chi}_{m,m}(z_1, z_2).$$

Now, according to the reasoning of Sect.1, the denominator should be a product including all factors of the form $(1 - t\, e(w_1)e(w_2))$ for all couples of nonzero weights w_1 and w_2 entering in the representations R_{λ_1} and R_{λ_2} of A_2. This gives

$$D_{diag}(t; z_1, z_2) = 1 + t^6 + (3 - z_1z_2)(t + t^5) + (6 + z_1^3 - 5z_1z_2 + z_2^3)(t^2 + t^4) + (7 + 2z_1^3 - 6z_1z_2 - z_1^2z_2^2 + 2z_2^3)t^3.$$

From this, we can find easily the generating function for the dimensions of the diagonal characters, which is

$$F_{diag}(t) = \frac{1 + 2t - 6t^2 + 2t^3 + t^4}{(1 - t)^6}$$

Thus, we try the ansatz

$$N_{diag}(t; z_1, z_2) = \sum_{l=0}^{4} C_l(z_1, z_2)t^l$$

for the numerator, and using the explicit form of the characters $\boldsymbol{\chi}_{m,m}(z_1, z_2)$ for $m = 0$ to 4, we arrive to a tentative form of $G_{diag}(t; z_1, z_2)$ as

$$G_{diag}(t; z_1, z_2) = \frac{1 + 2t - (z_1z_2 - 3)t^2 + 2t^3 + t^4}{D_{diag}(t; z_1, z_2)} \tag{20}$$

For this subset of characters, the operator Δ_t takes the form $\Delta_t = 3t^2\partial_t^2 + 9t\partial_t$ and one can check that $(\Delta_t - \Delta_z)G_{diag}(t; z_1, z_2) = 0$. Thus, (20) gives in fact the correct generating function.

The same procedure applied to C_2 gives the generating function of diagonal characters of that algebra. The result is

$$G_{diag}(t;z_1,z_2) = \frac{(1-t^2)(1+t^4+2tz_1+2t^3z_1+t^2(2z_1^2-z_1^2z_2+z_2+z_2^2))}{1+t^8-(t+t^7)d_1+(t^2+t^6)d_2+(t^3+t^5)d_3+t^4d_4}$$

where

$$\begin{aligned} d_1 &= z_1(-3+z_2) \\ d_2 &= -1+z_1^4+z_1^2(3-6z_2)+z_2+3z_2^2+z_2^3 \\ d_3 &= z_1(-3+2z_1^4-2z_2+8z_2^2+3z_2^3-z_1^2(-2+9z_2+z_2^2)) \\ d_4 &= z_1^6+z_1^4(4-6z_2)+z_1^2(-5-6z_2+5z_2^2)+z_2(-2+3z_2+4z_2^2+z_2^3). \end{aligned}$$

3 Some applications on weight multiplicities

The issue of the multiplicities of the weights entering in the irreducible representations of the simple Lie groups has been a subject of much research along the years [15]. In this section, we shall show how the above results can be used to obtain some generating functions for multiplicities in a very simple way. First, we describe the main points of the calculations by focusing in the case of zero weight multiplicities. Then, we give some other more general results for the case of the algebra C_2.

3.1 Generating function for zero weight multiplicities

The multiplicity $n_{m_1,m_2,\ldots,m_r}(p_1,p_2,\ldots,p_r)$ of the weight $p_1\lambda_1+p_2\lambda_2+\ldots+p_r\lambda_r$ in the representation $R_{m_1\lambda_1+m_2\lambda_2+\ldots+m_r\lambda_r}$ of the Lie algebra $\mathcal{A}$ can be computed as

$$\begin{aligned} n_{m_1,m_2,\ldots,m_r}(0,0,\ldots,0) &= \left(\frac{1}{2\pi}\right)^r \int_0^{2\pi} d\varphi_1 e^{-ip_1\varphi_1} \int_0^{2\pi} d\varphi_2 e^{-ip_2\varphi_2} \ldots \int_0^{2\pi} d\varphi_r e^{-ip_r\varphi_r} \boldsymbol{\chi}_{m_1,m_2,\ldots,m_r} \\ &= \left(\frac{1}{2\pi i}\right)^r \oint dx_1 \oint dx_2 \ldots \oint dx_r \frac{\boldsymbol{\chi}_{m_1,m_2,\ldots,m_r}}{x_1^{1+p_1}x_2^{1+p_2}\cdots x_r^{1+p_r}}\,; \end{aligned} \tag{21}$$

here the integrals in the second line are along the unit circles on the r complex planes parametrized by the complex coordinates $x_1,x_2,\ldots,x_r$. Then, in particular, the generating function for zero weight multiplicities

$$A_{0,0,\ldots,0}(t_1,t_2,\ldots,t_r) = \sum_{m_1=0}^{\infty}\sum_{m_2=0}^{\infty}\cdots\sum_{m_r=0}^{\infty} t_1^{m_1}t_2^{m_2}\cdots t_r^{m_r}\, n_{m_1,m_2,\ldots,m_r}(0,0,\ldots,0)$$

comes from the formula

$$A_{0,0,\ldots,0}(t_1,t_2,\ldots,t_r) = \left(\frac{1}{2\pi i}\right)^r \oint dx_1 \oint dx_2 \ldots \oint dx_r \frac{G(t_1,t_2,\ldots,t_r;z_1,z_2,\ldots,z_r)}{x_1x_2\cdots x_r},$$

which can be evaluated by means of the Cauchy's residue theorem. Let us see the results for the particular algebras that we have studied in the previous section.

3.1.1 The case of A_1

In this case, from (10) and (11), we have to compute

$$A_0(t_1) = \frac{1}{2\pi i}\oint dx_1 \frac{1}{x_1(1-t_1 z_1 + t_1^2)} = \frac{1}{2\pi i}\oint dx_1 \frac{x_1}{(t_1 - x_1)(t_1 x_1 - 1)}.$$

The convergence of the series requires $t_1 < 1$, while on the integration countour $|x_1| = 1$. So, the only pole of the integrand in the interior of the integration contour is a simple one at $x_1 = t_1$. Therefore

$$A_0(t_1) = \mathrm{Res}_{x_1 \to t_1}\left[\frac{x_1}{(t_1 - x_1)(t_1 x_1 - 1)}\right] = \frac{1}{1 - t_1^2}.$$

This is as it should be: the zero weight enter with multiplicity one in the representations $R_{m_1 \lambda_1}$ with m_1 even and does not appear in the represntations with m_1 odd.

3.1.2 The case of A_2

Now, we have to compute

$$A_{0,0}(t_1, t_2) = \left(\frac{1}{2\pi i}\right)^r \oint dx_1 \oint dx_2 \frac{G(t_1, t_2; z_1, z_2)}{x_1 x_2}$$

where, from (12), (13) and (16), the integrand is given by

$$\frac{G(t_1, t_2; z_1, z_2)}{x_1 x_2} = \frac{(1 - t_1 t_2) x_1 x_2}{(t_2 - x_1)(t_1 x_1 - 1)(t_1 - x_2)(t_2 x_1 - x_2)(t_1 x_2 - x_1)(t_2 x_2 - 1)}.$$

We perform the x_1 integral first. As $|x_1| = |x_2| = 1$ and $t_1, t_2 < 1$, there are poles inside the unit circle for $x_1 = t_2$ and $x_1 = t_1 x_2$. Thus

$$\begin{aligned} J_1(t_1, t_2, x_2) &= \frac{1}{2\pi i}\oint dx_1 \frac{G(t_1, t_2; z_1, z_2)}{x_1 x_2} = \{\mathrm{Res}_{x_1 \to t_2} + \mathrm{Res}_{x_1 \to t_1 x_2}\}\left[\frac{G(t_1, t_2; z_1, z_2)}{x_1 x_2}\right] \\ &= \frac{(1 + t_1 t_2) x_2}{(t_1 - x_2)(t_2^2 - x_2)(t_1^2 x_2 - 1)(t_2 x_2 - 1)}. \end{aligned}$$

As $J_1(t_1, t_2, x_2)$ has poles inside the x_2 unit circle at $x_2 = t_1$ and $x_2 = t_2^2$, we finally obtain the generating function for zero weight multiplicities as

$$\begin{aligned} A_{0,0}(t_1, t_2) &= \frac{1}{2\pi i}\oint dx_2 J_1(t_1, t_2, x_2) = \left\{\mathrm{Res}_{x_2 \to t_1} + \mathrm{Res}_{x_2 \to t_2^2}\right\} J_1(t_1, t_2, x_2) \\ &= \frac{1 - t_1^3 t_2^3}{(1 - t_1^3)(1 - t_1 t_2)^2 (1 - t_2^3)}. \end{aligned}$$

3.1.3 The case of C_2

For the Lie algebra C_2, according to (17), (18) and (19) the integrand takes the form

$$\frac{x_1^2 x_2((1+t_2)x_1x_2 + t_1^2 t_2(1+t_2)x_1x_2 - t_1t_2(1+x_2)(x_1^2+x_2))}{(x_1-t_1)(t_1x_1-1)(t_1x_1-x_2)(t_2x_1^2-x_2)(x_2-t_2)(x_1-t_1x_2)(x_1^2-t_2x_2)(t_2x_2-1)}.$$

The x_2 coordinate has poles with $|x_2| < 1$ at $x_2 = t_2$, $x_2 = t_1x_1$ and $x_2 = t_2x_1^2$. Applying the residue theorem, we find

$$\begin{aligned} J_2(t_1,t_2,x_1) &= \frac{1}{2\pi i}\oint dx_2 \frac{G(t_1,t_2;z_1,z_2)}{x_1x_2} = \\ &= -\frac{x_1(-(1+t_2^2)x_1 + t_1^2(t_2+t_2^3)x_1 + t_1(-1+t_2)t_2(1+x_1^2)}{(t_1^2-1)(t_2-1)(t_1-x_1)(t_2-x_1)(t_2+x_1)(t_1x_1-1)(t_2^2x_1^2-1)}. \end{aligned}$$

The poles of $J_2(t_1,t_2,x_1)$ into the x_1 unit circle are located at $x_1 = t_1$ and $x_1 = \pm t_2$. Thus, the final result is

$$A_{0,0}(t_1,t_2) = \frac{1}{2\pi i}\oint dx_1 J_2(t_1,t_2,x_1) = \frac{1+t_1^2t_2}{(1-t_1^2)^2(1-t_2)(1-t_2^2)}\,.$$

3.2 Some further results on weight multiplicities for C_2

Let us now specialize to the case of the algebra C_2 to study some other properties of the multiplicities of the weights entering in its irreducible representations. It will prove convenient to introduce a new generating function $H(t_1,t_2;y_1,y_2)$ defined as

$$H(t_1,t_2;y_1,y_2) = \sum_{m=0}^{\infty}\sum_{n=0}^{\infty}\sum_{p=0}^{\infty}\sum_{q=0}^{\infty} n_{p,q}(m,n)\, y_1^m y_2^n t_1^p t_2^q$$

where $n_{p,q}(m,n)$ is the multiplicity of the weight $m\lambda_1 + n\lambda_2$ in the representation $R_{p\lambda_1+q\lambda_2}$. From (21), one can compute $H(t_1,t_2;y_1,y_2)$ as a geometric series,

$$H(t_1,t_2;y_1,y_2) = \left(\frac{1}{2\pi i}\right)^2 \oint dx_1 \oint dx_2 \frac{G(t_1,t_2;z_1,z_2)}{(x_1-y_1)(x_2-y_2)},$$

and using (19) this gives

$$H(t_1,t_2;y_1,y_2) = \frac{a + b_1y_1 + b_2y_2 + c_{1,2}y_1y_2 + dy_1^2 + ey_1^2y_2}{(1-t_1^2)^2(1-t_2^2)(1-t_2)(1-t_1y_1)(1-t_2^2y_1^2)(1-t_1^2y_2)(1-t_2y_2)} \tag{22}$$

where

$$\begin{aligned} a &= 1+t_1^2t_2, & b_1 &= t_1t_2(1-t_1^2), & b_2 &= -t_1t_2(t_1^3+t_1t_2), \\ c_{1,2} &= t_1t_2(t_1^4-t_1^2), & d &= -t_1^2t_2^2(1+t_2), & e &= t_1^2t_2^2(t_1^2+t_1^2t_2+t_2^2-1). \end{aligned}$$

Now, we can rewrite $H(t_1, t_2; y_1, y_2)$ as a series in the variables y_1 and y_2

$$H(t_1, t_2; y_1, y_2) = \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} A_{m,n}(t_1, t_2) y_1^m y_2^n$$

such that the coefficient

$$A_{m,n}(t_1, t_2) = \sum_{p=0}^{\infty} \sum_{q=0}^{\infty} n_{p,q}(m, n) t_1^p t_2^q \tag{23}$$

is the generating function for the multiplicities of the weight $m\lambda_1 + n\lambda_2$ in all the irreducible representations of C_2. From (22), and after some tedious algebra, one can obtain the explicit form of these generating functions as

$$A_{m,n}(t_1, t_2) = \frac{t_1^{m+2n+2}(t_1^2 - t_2^2)(1 - t_2^2) - t_1^{m+2} t_2^{n+1}(1 - t_1^2)(1 - t_2^2) - t_2^{m+n+1}(t_1^2 - t_2)(1 - t_1^2) f(t_1, t_2)}{(1 - t_1^2)^2(1 - t_2^2)(1 - t_2)(t_1^2 - t_2^2)(t_1^2 - t_2)}$$

with

$$f(t_1, t_2) = \begin{cases} (t_1^2 + t_2) & \text{for } m \text{ even} \\ (1 + t_2) & \text{for } m \text{ odd} \end{cases}$$

thus generalizing the result for zero-weight multiplicities seen in the previous subsection.

In particular, one can use these expressions to write two recurrence relations among the multiplicities of the weights. The first one is a recurrence relation among the multiplicities of a fixed weight $m\lambda_1 + n\lambda_2$ in different representations. Let us call $X_{m,n}(t_1, t_2) = (1 - t_1^2)(1 - t_2)A_{m,n}(t_1, t_2)$. From the definition of $A_{m,n}(t_1, t_2)$, one has

$$X_{m,n}(t_1, t_2) = \sum_{p=0}^{\infty} \sum_{q=0}^{\infty} \left[n_{p,q}(m, n) - n_{p-2,q}(m, n) - n_{p,q-1}(m, n) + n_{p-2,q+1}(m, n) \right] t_1^p t_2^q$$

while the explicit expressions given above yield

$$\begin{aligned} X_{m,n}(t_1, t_2) &= \frac{(1 - t_2^2) \sum_{j=0}^{n-1} t_1^{m+2n-2j} t_2^j + (1 - t_2^2) \left[(1 + t_2) \sum_{j=0}^{\frac{m}{2}} t_1^{2j} t_2^{m+n-2j} - t_2^{m+n+1} \right]}{(1 - t_1^2)(1 - t_2^2)} \\ &+ \frac{t_1^2 t_2^{m+n+1} + t_1^{m+n+2}}{(1 - t_1^2)(1 - t_2^2)} \end{aligned}$$

for m even and

$$\begin{aligned} X_{m,n}(t_1, t_2) &= \frac{(1 - t_2^2) \sum_{j=0}^{n-1} t_1^{m+2n-2j} t_2^j + t_1(1 - t_2^2)(1 + t_2) \left[(1 + t_2) \sum_{j=0}^{\frac{m-3}{2}} t_1^{2j} t_2^{m+n-1-2j} + t_1^{m-1} t_2^n \right]}{(1 - t_1^2)(1 - t_2^2)} \\ &+ \frac{(1 + t_2) t_1 t_2^{m+n+1}}{(1 - t_1^2)(1 - t_2^2)} \end{aligned}$$

for m odd. Let us consider the formula for m even and compare it with the diagram of Fig. 1, which shows all representations $R_{p\lambda_1 + q\lambda_2}$ with nonzero multiplicity for the weight $m\lambda_1 + n\lambda_2$. The first term in $X_{m,n}(t_1,t_2)$ can be expanded as a geometric series which contains, always with coefficient equal to one, all the products $t_1^p t_2^q$ for p and q corresponding to points in the diagonals beginning in the segment AB. In a similar way, the second term in $X_{m,n}(t_1,t_2)$ gives all such products for the points in the diagonals normal to the line from (m,n) to C, the third corresponds to the diagonals beginning in $D', E', F', \ldots$, etc, and the fourth to the diagonals from $D, E, F, \ldots$ etc. From this and an analogous analysis for m odd, we can finally conclude that

$$n_{p,q}(m,n) - n_{p-2,q}(m,n) - n_{p,q-1}(m,n) + n_{p-2,q+1}(m,n) = y_{p,q}(m,n) \tag{24}$$

where $y_{p,q}(m,n) = 1$ if (p,q) labels a irreducible representations of C_2 containing the weight $m\lambda_1 + n\lambda_2$ (except for m even, $p = 0$ and q of opposite parity to n which gives $y_{p,q}(m,n) = 0$) and $y_{p,q}(m,n) = 0$ if the weight $m\lambda_1 + n\lambda_2$ is not in $R_{p\lambda_1+q\lambda_2}$.

There is a second recurrence relation, this time among the multiplicities of weights of the same representation. From the previous expressions of $A_{m,n}(t_1,t_2)$, one can obtain

$$R_{m,n}(t_1,t_2) = A_{m,n}(t_1,t_2) - A_{m+2,n}(t_1,t_2) - A_{m,n+1}(t_1,t_2) + A_{m+2,n+1}(t_1,t_2) \tag{25}$$

as

$$R_{m,n}(t_1,t_2) = \frac{1}{1-t_2}\sum_{j=0}^{n} t_1^{m+2n-2j} t_2^j + \frac{1}{1-t_2}\sum_{j=0}^{\frac{m}{2}-1} t_1^{2j} t_2^{m+n-2j} - \frac{t_2^{m+n+1}}{(1-t_1^2)(1-t_2)}$$

for m even and

$$R_{m,n}(t_1,t_2) = \frac{1}{1-t_2}\sum_{j=0}^{n} t_1^{m+2n-2j} t_2^j + \frac{1}{1-t_2}\sum_{j=0}^{\frac{m}{2}-\frac{3}{2}} t_1^{2j+1} t_2^{m+n-1-2j} - \frac{t_1 t_2^{m+n+1}}{(1-t_1^2)(1-t_2)}$$

for m odd. Expanding the denominators as geometric series and using (23), one finds that in both cases

$$n_{p,q}(m,n) - n_{p,q}(m+2,n) - n_{p,q}(m,n+1) + n_{p,q}(m+2,n+1) = \varepsilon_{p,q}(m,n) \tag{26}$$

where the right-hand member is a sum of three terms, $\varepsilon_{p,q}(m,n) = X + Y - Z$, which are zero except for the cases

$$\begin{array}{rll} X = 1 & \text{when} & m \le p \ \text{ and } \ p \le m+2n \le p+2q \\ Y = 1 & \text{when} & m \ge p+2 \ \text{ and } \ m+n \le p+q \\ Z = 1 & \text{when} & m+n \le q-1. \end{array}$$



Figure 1. The highest weights (modulo Weyl reflections) corresponding to representations containing the weight $m\lambda_1 + n\lambda_2$. Here, λ_1 is diagonal of magnitude one and λ_2 is vertical of magnitude $\sqrt{2}$.

As $X = 1$ and $Y = 1$ do not occur at the same time, $\varepsilon_{p,q}(m,n)$ is always 1, 0 or -1. To describe the domains $D1$, $D0$ and $D(-1)$ in which $\epsilon_{p,q}(m,n)$ takes these values, let us consider the weight diagram of the representation $p\lambda_1 + q\lambda_2$ using now Cartesian coordinates (x,y) (with $x = m$, $y = m + 2n$) instead of the m and n labels of the weights. The weights entering in the diagram form a square lattice with a spacing of one unit which includes the highest weight of the representation and is contained in the polygon of vertices $(0, p+2q)$, $(p, p+2q)$, $(p+q, p+q)$ and $(0,0)$. If we call R_X, R_Y and R_Z the regions of the diagram in which, respectively, $X = 1$, $Y = 1$ and $Z = 1$ it follows that:

- If $2q-2 < p$, $R_X \cup R_Y$ does not intersect R_Z. Thus, $D1 = R_X \cup R_Y$, $D(-1) = R_Z$, and $D0$ the remaining weights. So, in this case $D1$ is the upper region of the diagram, starting from the horizontal line $y = p$, $D0$ is the area below that line and above the diagonal $y = 2q - 2 - x$, and $D(-1)$ includes that diagonal and the weights below it, see fig. 2.
- If $p \leq 2q - 2 \leq 2p$ the intersection $R_X \cap R_Z$ is the (possibly degenerate) triangle T of vertices $(0, 2q-2)$, $(0, p)$ and $(2q-2-p, p)$. In this case $D1 = R_X \cup R_Y - T$, $D(-1) = R_Z - T$, and





Figure 2. The weights of the representation $R_{10\lambda_1+5\lambda_2}$ in the domains $D1$, $D0$ and $D(-1)$ are marked, respectively, with black dots, circles and encircled black dots. The region $R_X \cup R_Y$ has perimeter $ABCDEA$, while R_Z is contained in $FGOF$. The number over a weight means its multiplicity.

Figure 3. Weights of the representation $R_{10\lambda_1+9\lambda_2}$. The region $R_X \cup R_Y$ is bounded by $AFBCDEGA$, R_Z is into $FGHOAF$ and the triangle T is $FGAF$.

$D0$ the remaining weights. Now, as one can see in fig. 3, some weights above $y = p$ located near the y axis are in D0 instead of in D1.

- If $2q-2 > 2p$ the intersection $(R_X \cup R_Y) \cap R_Z$ is the cuadrilateral K of vertices $(0,p)$, $(0,2q-2)$, $(q-1,q-1)$ and (p,p). Therefore $D1 = R_X \cup R_Y - C$, $D(-1) = R_Z - C$, and $D0$ the remaining weights, see fig. 4.





Figure 4. Weights of the representation $R_{5\lambda_1+10\lambda_2}$. The region $R_X \cup R_Y$ is bounded by $AFBCDGEAF$, R_Z is into $FGEOAF$ and the cuadrilateral K is $FGEAF$.

Figure 5. The multiplicity of $m\lambda_1 + n\lambda_2$ in the representation $R_{p\lambda_1+q\lambda_2}$ is given by the number of weights marked with a square. By parity, the weight signaled with X is excluded.

These recurrence relations provide useful information on the multiplicities and, in fact, can be used to devise some simple rules to compute the multiplicity of any desired weight on a given representation. As an example, let us take the case of m even and consider a situation with (m, n) and (p, q) as given in the figure 5. We can write the first recurrence relation as

$$n_{p,q}(m,n) - n_{p,q-1}(m,n) = y_{p,q}(m,n) + n_{p-2,q}(m,n) - n_{p-2,q-1}(m,n)$$

and iterating we find

$$\begin{aligned} n_{p,q}(m,n) &= y_{p,q}(m,n) + y_{p-2,q}(m,n) + \cdots + y_{2,q}(m,n) + y_{0,q}(m,n) \\ &+ y_{p,q-1}(m,n) + y_{p-2,q-1}(m,n) + \cdots + y_{2,q-1}(m,n) + y_{0,q-1}(m,n) \\ &+ y_{p,q-2}(m,n) + y_{p-2,q-2}(m,n) + \cdots + y_{2,q-2}(m,n) + y_{0,q-2}(m,n) + \cdots \end{aligned}$$

so that finally

$$n_{p,q}(m,n) = \sum_{\mu \in R} y_{\mu}(m,n)$$

where the sum is over to the set R of weights marked in the figure. Thus, $n_{p,q}(m,n)$ can be obtained by simply counting the number of poits in R except those in the vertical axis with opposite parity of q and n, which have $y_\mu(m,n)=0$. This rule can be used to obtain some explicit formulae. For instance, for the case p and q even and $q\leq\frac{p}{2}$, one finds that the multiplicities of the weights on borders of the diagram are given by

$$\begin{aligned} n_{p,q}(p+q-2s,0) &= (s+1)^2 && \text{for} \quad 0\leq s\leq \frac{q}{2} \\ n_{p,q}(p-2s,0) &= (\frac{q}{2}+1)^2+s(q+1) && \text{for} \quad 0\leq s\leq \frac{p-q}{2} \\ n_{p,q}(2s,0) &= n_{p,q}(0,0)-s^2 && \text{for} \quad 0\leq s\leq \frac{q}{2} \end{aligned}$$

and

$$\begin{aligned} n_{p,q}(0,\frac{p}{2}+q-s) &= \frac{(s+1)(s+2)}{2} && \text{for} \quad 0\leq s\leq q \\ n_{p,q}(0,\frac{p}{2}-s) &= \frac{(s+1)(s+2)}{2}+s(q+1) && \text{for} \quad 0\leq s\leq \frac{p}{2}-q \\ n_{p,q}(0,s)-n_{p,q}(0,s+1) &= s+1-\theta(s+1) && \text{for} \quad 0\leq s\leq q-1 \end{aligned}$$

where $\theta(r)$ is one (zero) for r even (odd). One can then use (26) to compute the multiplicities of the inner weights.

Appendix

In this Appendix, we present the Calogero-Sutherland differential operator whose eigenfunctions are the characters of the algebra C_2 and give a short collection of characters of C_2 which includes those needed to fix the numerator of the generating function $G(t_1,t_2;z_1,z_2)$. For a more detailed account of the use of the quantum Calogero-Sutherland models to deal with the characters or other generalized orthogonal polynomials related to Lie algebras, see [13] and references therein.

The general form of the differential operator we are looking for is

$$\Delta_z = a_{1,0}(z_1,z_2)\partial_{z_1}^2+a_{0,1}(z_1,z_2)\partial_{z_2}^2+a_{1,1}(z_1,z_2)\partial_{z_1}\partial_{z_2}+b_1(z_1,z_2)\partial_{z_1}+b_2(z_1,z_2)\partial_{z_2}$$

and we will fix the coefficients using that the character $\boldsymbol{\chi}_{m_1,m_2}$ of the irreducible representation of C_2 with highest weight $\lambda=m_1\lambda_1+m_2\lambda_2$ is an eigenfunction of the Schrödinger equation

$$\Delta_z\boldsymbol{\chi}_{m_1,m_2}=\varepsilon(m_1,m_2)\boldsymbol{\chi}_{m_1,m_2}$$

with eigenvalue

$$\varepsilon(m_1,m_2)=\langle\lambda,\lambda\rangle+2\langle\lambda,\rho\rangle,$$

where $\langle\cdot,\cdot\rangle$ is the scalar product in the space spanned by the orthonormal basis $\{e_1,e_2\}$, the two fundamental weights of C_2 are represented in this basis by

$$\lambda_1=e_1, \qquad \lambda_2=e_1+e_2,$$

the four positive roots are

$$\alpha_1 = e_1 - e_2, \qquad \alpha_2 = 2e_2, \qquad \alpha_3 = e_1 + e_2, \qquad \alpha_4 = 2e_1,$$

and $\rho = \lambda_1 + \lambda_2$ is the Weyl vector of the algebra [16, 17], so that

$$\varepsilon(m_1, m_2) = m_1^2 + 2m_2^2 + 2m_1 m_2 + 4m_1 + 6m_2.$$

On the other hand, given that

$$\begin{aligned} \boldsymbol{\chi}_{1,0}(z_1, z_2) &= z_1 \\ \boldsymbol{\chi}_{0,1}(z_1, z_2) &= z_2, \end{aligned}$$

the direct products of the representations of C_2 [16, 17]

$$\begin{aligned} R_{1,0} \otimes R_{1,0} &= R_{2,0} + R_{0,1} + R_{0,0} \\ R_{1,0} \otimes R_{0,1} &= R_{1,1} + R_{1,0} \\ R_{0,1} \otimes R_{0,1} &= R_{0,2} + R_{2,0} + R_{0,0} \end{aligned}$$

allow us to solve for

$$\begin{aligned} \boldsymbol{\chi}_{2,0}(z_1, z_2) &= z_1^2 - z_2 - 1 \\ \boldsymbol{\chi}_{1,1}(z_1, z_2) &= z_1 z_2 - z_1 \\ \boldsymbol{\chi}_{0,2}(z_1, z_2) &= z_2^2 - z_1^2 + z_2. \end{aligned}$$

Using all these characters in the Schrödinger equation one finds

$$\begin{aligned} a_{1,0}(z_1, z_2) &= z_1^2 - 2z_2 - 6 \\ a_{0,1}(z_1, z_2) &= 2z_2^2 - 4z_1^2 + 4z_2 - 6 \\ a_{1,1}(z_1, z_2) &= 2z_1 z_2 - 10z_1 \\ b_1(z_1, z_2) &= 5z_1 \\ b_{(}z_1, z_2) &= 4z_2. \end{aligned}$$

Once the operator Δ_z is known, other characters of C_2 can be computed by solving the Schrödinger equation or by expansion of generating function (19). We give here a few of them:

$$\begin{aligned} \boldsymbol{\chi}_{0,3}(z_1, z_2) &= -1 + z_1^2 - 2z_1^2 z_2 + 2z_2^2 + z_2^3 \\ \boldsymbol{\chi}_{1,2}(z_1, z_2) &= z_1 - z_1^3 + z_1 z_2^2 \\ \boldsymbol{\chi}_{2,1}(z_1, z_2) &= 1 - z_1^2 - z_2 + z_1^2 z_2 - z_2^2 \\ \boldsymbol{\chi}_{3,0}(z_1, z_2) &= -z_1 + z_1^3 - 2z_1 z_2 \\ \boldsymbol{\chi}_{0,4}(z_1, z_2) &= -z_1^2 + z_1^4 - 2z_2 + z_2^2 - 3z_1^2 z_2^2 + 3z_2^3 + z_2^4 \\ \boldsymbol{\chi}_{1,3}(z_1, z_2) &= -2z_1 + 2z_1^3 - 2z_1^3 z_2 + z_1 z_2^2 + z_1 z_2^3 \\ \boldsymbol{\chi}_{2,2}(z_1, z_2) &= 2z_1^2 - z_1^4 + z_1^2 z_2 - 2z_2^2 + z_1^2 z_2^2 - z_2^3 \end{aligned}$$

$$
\begin{aligned}
\boldsymbol{\chi}_{3,1}(z_1,z_2) &= 2z_1 - z_1^3 + z_1^3 z_2 - 2z_1 z_2^2 \\
\boldsymbol{\chi}_{4,0}(z_1,z_2) &= -z_1^2 + z_1^4 + 2z_2 - 3z_1^2 z_2 + z_2^2 \\
\boldsymbol{\chi}_{0,5}(z_1,z_2) &= 3z_1^2 - 2z_1^4 - 2z_2 + 3z_1^4 z_2 - 3z_2^2 - 3z_1^2 z_2^2 + 3z_2^3 - 4z_1^2 z_2^3 + 4z_2^4 + z_2^5 \\
\boldsymbol{\chi}_{1,4}(z_1,z_2) &= 2z_1 - 3z_1^3 + z_1^5 - 2z_1 z_2 + 2z_1^3 z_2 - 3z_1^3 z_2^2 + 2z_1 z_2^3 + z_1 z_2^4 \\
\boldsymbol{\chi}_{2,3}(z_1,z_2) &= 1 - 4z_1^2 + 2z_1^4 + 2z_2 + z_1^2 z_2 - 2z_1^4 z_2 - z_2^2 + 3z_1^2 z_2^2 - 3z_2^3 + z_1^2 z_2^3 - z_2^4 \\
\boldsymbol{\chi}_{3,2}(z_1,z_2) &= -z_1 + 2z_1^3 - z_1^5 + 2z_1^3 z_2 - 2z_1 z_2^2 + z_1^3 z_2^2 - 2z_1 z_2^3 \\
\boldsymbol{\chi}_{4,1}(z_1,z_2) &= -1 + 2z_1^2 - z_1^4 - z_2 + z_1^2 z_2 + z_1^4 z_2 + 2z_2^2 - 3z_1^2 z_2^2 + z_2^3 \\
\boldsymbol{\chi}_{5,0}(z_1,z_2) &= -z_1 - z_1^3 + z_1^5 + 4z_1 z_2 - 4z_1^3 z_2 + 3z_1 z_2^2 \\
\boldsymbol{\chi}_{0,6}(z_1,z_2) &= 1 - 3z_1^2 + 3z_1^4 - z_1^6 + 6z_1^2 z_2 - 3z_1^4 z_2 - 6z_2^2 + 6z_1^4 z_2^2 - 3z_2^3 - 8z_1^2 z_2^3 + 6z_2^4 - 5z_1^2 z_2^4 \\
&+ 5z_2^5 + z_2^6 \\
\boldsymbol{\chi}_{1,5}(z_1,z_2) &= -2z_1 + 6z_1^3 - 3z_1^5 - 2z_1^3 z_2 + 3z_1^5 z_2 - 3z_1 z_2^2 + z_1 z_2^3 - 4z_1^3 z_2^3 + 3z_1 z_2^4 + z_1 z_2^5 \\
\boldsymbol{\chi}_{2,4}(z_1,z_2) &= -1 + 4z_1^2 - 4z_1^4 + z_1^6 + 2z_2 - 3z_1^2 z_2 + z_1^4 z_2 + 3z_2^2 + 3z_1^2 z_2^2 - 3z_1^4 z_2^2 - 3z_2^3 \\
&+ 5z_1^2 z_2^3 - 4z_2^4 + z_1^2 z_2^4 - z_2^5 \\
\boldsymbol{\chi}_{3,3}(z_1,z_2) &= 2z_1 - 5z_1^3 + 2z_1^5 + 4z_1 z_2 - z_1^3 z_2 - 2z_1^5 z_2 + 5z_1^3 z_2^2 - 4z_1 z_2^3 + z_1^3 z_2^3 - 2z_1 z_2^4 \\
\boldsymbol{\chi}_{4,2}(z_1,z_2) &= -z_1^2 + 2z_1^4 - z_1^6 - z_2 - 3z_1^2 z_2 + 3z_1^4 z_2 + z_2^2 - 3z_1^2 z_2^2 + z_1^4 z_2^2 + 3z_2^3 - 3z_1^2 z_2^3 + z_2^4 \\
\boldsymbol{\chi}_{5,1}(z_1,z_2) &= -z_1 + 2z_1^3 - z_1^5 - 5z_1 z_2 + 2z_1^3 z_2 + z_1^5 z_2 + 3z_1 z_2^2 - 4z_1^3 z_2^2 + 3z_1 z_2^3 \\
\boldsymbol{\chi}_{6,0}(z_1,z_2) &= 1 - 2z_1^2 - z_1^4 + z_1^6 - z_2 + 6z_1^2 z_2 - 5z_1^4 z_2 - 3z_2^2 + 6z_1^2 z_2^2 - z_2^3
\end{aligned}
$$

Acknowledgement

J. F. N. acknowledges financial support from MTM2009-11154 project, SGPI-DGI(MICINN), Spain.

References

[1] I.G. Macdonald, *Symmetric Functions and Hall Polynomials*, Oxford, 1995.

[2] H. Weyl, Math. Zs. 23, 271–309 (1925); 24, 328–376, 377–395 (1926).

[3] H. Weyl, *Classical Groups. Their Invariants and Representations.* Princeton Univ. Press, 1946.

[4] W. García Fuertes and A.M. Perelomov, J. Phys. A: Math. Gen **35**, L335–L340 (2002).

[5] N. Okeke and M.A.Walton, J. Phys. A: Math. Theor **40**, 8873-8901 (2007).

[6] F. Calogero, J. Math. Phys. **12**, 419–436 (1971).

[7] B. Sutherland, Phys. Rev. **A4**, 2019–2021 (1972).

[8] J. Moser, Adv. Math. **16**, 197–220 (1975).

[9] M.A. Olshanetsky and A.M. Perelomov, Invent. Math. **37**, 93–108 (1976).

[10] M.A. Olshanetsky and A.M. Perelomov, Phys. Rep. **71**, 314–400, 1981; Phys. Rep. **94**, 313–404 (1983).

[11] A.M. Perelomov, J. Phys. **A31**, L31–L37 (1998); J. Phys. **A32**, 8563–8576 (1999).

[12] A.M. Perelomov, E. Ragoucy and Ph. Zaugg, J. Phys. **A31**, L559–L565 (1998).

[13] J. Fernández Núñez, W. García Fuertes, A.M. Perelomov, Phys. Lett. **A307**, 233–238 (2003); J. Math. Phys. **44**, 4957–4974(2003); J. Nonlinear Math. Phys. **12**, Suppl. 1, 280-301(2005); J. Math. Phys. **46**, 073508(2005); J. Math. Phys. **46**, 103505(2005); Theor. Math. Phys., **154**, 240-249(2008).

[14] A.M. Perelomov, *Proc. Clausthal Conf. 1999:Lie Theory and Its Applications in Physics III*, 139–154, World Scientific, 2000.

[15] E.P. Wigner, J. Phys. Rev. 51, 106–119 (1937).
H. Freudenthal, Proc. Nederl. Akad. Wet., A57, 369-376, 487-491 (1954).
G. Racah, *Lectures on Lie groups*, in *Group theoretical concepts and methods in elementary particle physics* (Lectures Istanbul Summer School Theoret. Phys., 1962), pages 1-36. Gordon and Breach, New York (1964).
B. Kostant, Trans. Amer. Math. Soc. 93, 53-73 (1959).
R.V. Moody and J. Patera, Bull. Amer. Math. Soc. 7, 237-242 (1982).

[16] A.L. Onishchik and E.B. Vinberg, *Lie Groups and Algebraic Groups,* Springer, 1990.

[17] Cornwell J.F., *Group Theory in Physics, vol II,* Academic Press, 1984; Bourbaki N. *Groupes et Algebres de Lie,* Hermann, 1975.